


Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Nowhere-zero 4-flow in almost Petersen-minor free graphs

Xiaofeng Wang^a, Cun-Quan Zhang^{a,*}, Taoye Zhang^b

^a Department of Mathematics, West Virginia University, Morgantown, WV 26506-6310, United States

^b Department of Mathematics, Pennsylvania State University, Dunmore, PA 18512, United States

Received 22 February 2006; accepted 29 November 2007

Available online 31 December 2007

Abstract

Tutte [W.T. Tutte, On the algebraic theory of graph colorings, J. Combin. Theory 1 (1966) 15–20] conjectured that every bridgeless Petersen-minor free graph admits a nowhere-zero 4-flow. Let $(P_{10})_{\bar{\mu}}$ be the graph obtained from the Petersen graph by contracting μ edges from a perfect matching. In this paper we prove that every bridgeless $(P_{10})_{\bar{3}}$ -minor free graph admits a nowhere-zero 4-flow.

© 2007 Elsevier B.V. All rights reserved.

Keywords: Integer flow; 4-flow; Edge-3-coloring; Petersen minor; Almost Petersen-minor free graphs

1. Introduction

The concept of integer flow was introduced by Tutte as a generalization of map coloring problem. The following conjecture is one of the major open problems in graph theory.

Conjecture 1.1 (Tutte [14]). *Every bridgeless graph without a Petersen minor admits a nowhere-zero 4-flow.*

For planar graphs, admitting a nowhere-zero 4-flow is equivalent to having a face 4-coloring. Hence, by the 4-Color Theorem [1–3,10], Conjecture 1.1 has been verified for all planar graphs. Furthermore, it was also announced that Conjecture 1.1 was verified for all cubic graphs [11,12]. By the Kuratowski Theorem, a graph is planar if and only if it contains neither K_5 -minor nor $K_{3,3}$ -minor. By applying the 4-Color Theorem, Conjecture 1.1 was further verified for $K_{3,3}$ -minor free graphs [15], K_5 -minor free graphs [7], and P_{10}^- -minor free graphs [13]. Each of these families contains the family of all planar graphs and may not necessarily be cubic. Graphs K_5 , $K_{3,3}$, P_{10} and P_{10}^- are illustrated in Figs. 1–5.

Let P_{10} be the Petersen graph with the exterior pentagon $1'2'3'4'5'1'$, interior pentagon $1''3''5''2''4''1''$ and a perfect matching $M = \{e_i = i'i'' : i = 1, 2, 3, 4, 5\}$. Let $(P_{10})_{\bar{\mu}}$ be the graph obtained from P_{10} by contracting F , where $F \subseteq M$ and $|F| = \mu$.

* Corresponding author.

E-mail address: cqzhang@math.wvu.edu (C.-Q. Zhang).




Fig. 1.




Fig. 2.




Fig. 3.

Remark. It is not hard to see that if M and M' are two perfect matchings of P_{10} , $F \subseteq M$, $F' \subseteq M'$ and $|F| = |F'|$, then $P_{10}/F \cong P_{10}/F'$. Hence $(P_{10})_{\bar{\mu}}$ is well defined.

The following is our main theorem.

Theorem 1.2. *Let G be a bridgeless graph. If G does not have a $(P_{10})_{\bar{3}}$ -minor, then G admits a nowhere-zero 4-flow.*

2. Notation and terminologies

For terms that are not defined here, readers can refer to textbooks [4,8,16] (for flows).

Let $G = (V, E)$ be a graph with vertex set V and edge set E and let D be an orientation of G . For a vertex $v \in V(G)$, let $E^+(v)$ (or $E^-(v)$) be the set of all arcs of $D(G)$ with their tails (or heads, respectively) at the vertex v . G is said to admit a nowhere-zero k -flow if there exists an ordered pair (D, f) , where $f : E(G) \rightarrow \{\pm 1, \pm 2, \dots, \pm(k-1)\}$, such that

$$\sum_{e \in E^+(v)} f(e) = \sum_{e \in E^-(v)} f(e)$$

for every vertex $v \in V(G)$. A graph G is a 4-flow snark if it is bridgeless and does not admit a nowhere-zero 4-flow. Let G and H be two graphs. If G contains a subgraph which is contractible to H , then H is a minor of G and we say G contains an H -minor. A 4-flow snark G is minor-prime if no proper minor of G is a 4-flow snark. With the definitions above, Conjecture 1.1 can be restated as follows.

Conjecture 2.1. *The Petersen graph is the only minor-prime 4-flow snark.*

Let H be a minor of a connected graph G . Then there is an onto mapping $f : V(G) \mapsto V(H)$ such that $f^{-1}(v)$ induces a connected subgraph $G[f^{-1}(v)]$ of G for every $v \in V(H)$ and H can be obtained from a spanning subgraph of G by contracting the edges of $G[f^{-1}(v)]$ for all $v \in V(H)$. Here f is called a minor mapping and $f^{-1}(v)$ is called a v -domain of f . A k -separator of a graph G is an ordered triple $(H_1, H_2; T)$ such that $H_1 \cup H_2 = G$ and $V(H_1 \cap H_2) = T$, where T is a vertex subset of G and $|T| = k$. Sometimes we say T is a k -separator if there is no




Fig. 4.




Fig. 5.




Fig. 6.

confusion. A k -separator $(H_1, H_2; T)$ of G is *trivial* if one of H_1 and H_2 , say H_1 , is acyclic. G is *quasi- k -connected* if G is 3-connected and every t -separator of G with $t \leq k$ is trivial. Let x be a vertex of G . The vertex x *separates* G into q parts H_1, \dots, H_q if $G = H_1 \cup \dots \cup H_q$ and $V(H_i \cap H_j) = \{x\}$ for every pair of $i \neq j$.

Let X be a connected subgraph of G and Y_1, Y_2, Y_3, Y_4 be four disjoint connected subgraphs of $G - V(X)$ and $X \cap N(Y_i) \neq \emptyset$ for $i = 1, 2, 3, 4$ where $N(Y_i)$ denotes the set of neighbors of Y_i . Let $J = \{Y_1, Y_2, Y_3, Y_4\}$. For each 2×2 -partition $P = \{\{a, b\}, \{c, d\}\}$ of $\{1, 2, 3, 4\}$, X is P -splittable if X contains two disjoint paths Q and Q' such that Q joins $X \cap N(Y_a)$ and $X \cap N(Y_b)$, Q' joins $X \cap N(Y_c)$ and $X \cap N(Y_d)$, $i \in \{a, b, c, d\}$. An example of a $\{\{1, 2\}, \{3, 4\}\}$ -splittable subgraph is illustrated in Fig. 6. X is k -splittable with respect to J if there are k distinct 2×2 partitions P_1, \dots, P_k of $\{1, 2, 3, 4\}$ such that X is P_i -splittable for each $i = 1, \dots, k$. (Remark: $k \leq 3$.) An example of a 2-splittable subgraph is illustrated in Fig. 7.

3. Lemmas

Lemma 3.1 (Catlin [5]). *If G is a minor-prime 4-flow snark, then the girth of G is at least 5.*

Lemma 3.2 (Lai, Li and Poon [7]). *If a bridgeless graph G does not admit a nowhere-zero 4-flow, then G has a K_5 -minor.*




Fig. 7.

Lemma 3.3 (Thomas and Thomson, Lemma 4.4 of [13]). *If G is a minor-prime 4-flow snark, then G is quasi-4-connected (that is, every k -separator of G is trivial for each $k \leq 3$).*

Obviously, Lemma 3.3 generalizes Lemma 3.1 and Theorem 3.7.15 of [16].

Proposition 3.4. *Let X be a connected subgraph of G and Y_1, Y_2, Y_3, Y_4 be four disjoint connected subgraphs of $G - V(X)$ where $V(X) \cap N(Y_i) \neq \emptyset$ for $i = 1, 2, 3, 4$. Let k be the greatest integer such that X is k -splittable with respect to $J = \{Y_1, Y_2, Y_3, Y_4\}$.*

- (i) *If $k \leq 1$, say, X is $\{\{1,2\},\{3,4\}\}$ -splittable or 0-splittable, then X has a 1-separator $(H_1, H_2; \{x\})$ such that $V(X) \cap [N(Y_1) \cup N(Y_2)] \subseteq V(H_1)$ and $V(X) \cap [N(Y_3) \cup N(Y_4)] \subseteq V(H_2)$.*
- (ii) *If $k = 0$, then there exists a cut vertex x of X that separates X into four parts H_1, H_2, H_3, H_4 such that $V(X) \cap N(Y_i) \subseteq V(H_i)$ for each i .*

Proof. (i) Let G_1 be the graph induced by $X \cup Y_1 \cup Y_2 \cup Y_3 \cup Y_4$. Let G_2 be the graph obtained from G_1 by contracting each Y_i into a single vertex y_i for $i = 1, 2, 3, 4$, and deleting all edges between y_i and y_j for all $\{i, j\} \subset \{1, 2, 3, 4\}$. Note that G_2 is connected since $V(X) \cap N(Y_i) \neq \emptyset$ for $i = 1, 2, 3, 4$.

Since X is neither $\{\{1,3\},\{2,4\}\}$ -splittable nor $\{\{1,4\},\{2,3\}\}$ -splittable, it is impossible that there is a pair of disjoint paths joining $\{y_1, y_2\}$ and $\{y_3, y_4\}$. By Menger's theorem, there is a cut vertex $x \in V(G_2)$ that separates $\{y_1, y_2\}$ and $\{y_3, y_4\}$. It is obvious that $x \in V(X)$. That is, X has a 1-separator $(H_1, H_2; x)$ that $N_{G_2}(y_1) \cup N_{G_2}(y_2) \subseteq V(H_1)$ and $N_{G_2}(y_3) \cup N_{G_2}(y_4) \subseteq V(H_2)$.

(ii) Continue from (i). Assume that there is a path P_1 joining y_1 and y_2 in the graph $G_2 - \{x\}$ (without passing through x). Note that x is a cut vertex that separates $\{y_1, y_2\}$ and $\{y_3, y_4\}$. Thus, this path P_1 is contained in the induced subgraph $G_2[V(H_1 - x) \cup \{y_1, y_2\}]$ and there is another path P_2 joining y_3 and y_4 in the induced subgraph $G_2[H_2 \cup \{y_3, y_4\}]$ since H_2 is connected. This contradicts that X is 0-splittable. So every path from y_1 to y_2 must go through x . Symmetrically, every path from y_3 to y_4 must go through x as well. That implies each component of $X - x$ is adjacent to at most one of $\{y_1, y_2, y_3, y_4\}$. ■

4. Proof of the main theorem

Let G be a minor-prime 4-flow snark. By Lemma 3.3, G is quasi-4-connected. By Lemma 3.2, K_5 is a minor of G . Let $V(K_5) = \{v_1, v_2, v_3, v_4, v_5\}$, and $f : V(G) \mapsto V(K_5)$ be a minor mapping.

If G does not contain a $(P_{10})_4$ -minor, then v_a -domain $f^{-1}(v_a)$ is at most 0-splittable with respect to $\{f^{-1}(v_{i_j}) : j = 1, 2, 3, 4\}$ for every $\{a, i_1, i_2, i_3, i_4\} = \{1, 2, 3, 4, 5\}$. By Proposition 3.4(ii), each $f^{-1}(v_a)$ has a cut vertex v_a^* that separates $N(f^{-1}(v_{i_j}))$ for $j = 1, 2, 3, 4$. Hence $\{v_i^*, v_j^*\}$ is a 2-separator of G . Since $G \neq K_5$, there exist $\{i, j\} \subseteq \{1, 2, 3, 4, 5\}$ such that $\{v_i^*, v_j^*\}$ is a non-trivial 2-separator. This contradicts the fact that G is quasi-4-connected. Hence G contains $(P_{10})_4$ as a minor.¹

Let $f : V(G) \rightarrow (P_{10})_4$ be a minor mapping where the vertex set of $(P_{10})_4$ is $\{v_{1'}, v_{1''}, v_2, v_3, v_4, v_5\}$, the contraction of the edge $v_{1'}v_{1''}$ yields a K_5 , $v_{1'}$ is adjacent to v_2 and v_5 , and $v_{1''}$ is adjacent to v_3 and v_4 . Let $U_i = f^{-1}(v_i)$ for $i \in \{1', 1'', 2, 3, 4, 5\}$ (see Fig. 8). Define $U_1 = U_{1'} \cup U_{1''}$ and choose a minor mapping f such that $|U_1|$ is as small as possible. Now assume that G does not contain a $(P_{10})_3$ -minor.

¹ It was suggested by a referee that this part of the proof can be obtained directly by applying the Splitter Theorem (see [6,9]). Here, for the purpose of completeness, we include this short proof.




Fig. 8.




Fig. 9.

Claim 1. $|U_{1'}| = |U_{1''}| = 1$.

Proof. It is clear that $|U_{1'}| \geq 1$ and $|U_{1''}| \geq 1$.

Let $e = u_{1'}u_{1''}$ be an edge between $U_{1'}$ and $U_{1''}$ where $u_{1'} \in U_{1'}$, $u_{1''} \in U_{1''}$. Since each of $U_{1'}$ and $U_{1''}$ is connected, there are spanning trees T_1 and T_2 of $U_{1'}$ and $U_{1''}$, respectively. Let $T = T_1 \cup T_2 \cup \{e\}$. T is a spanning tree of U_1 .

Assume there exist $w_2 \in N(U_2) \cap U_{1'}$ and $w_5 \in N(U_5) \cap U_{1'}$ such that $w_2 \neq w_5$. Since T_1 is a spanning tree of $U_{1'}$, there is a unique path P_2 from $u_{1'}$ to w_2 in T_1 , and a path P_5 from $u_{1'}$ to w_5 in T_1 . Without loss of generality, we may assume that P_2 is not shorter than P_5 . Since $w_2 \neq w_5$, P_5 does not contain w_2 . Let C_2 be the set of vertices of the component of $T_1 \setminus P_5$ that contains w_2 . Now we define a new minor mapping f_1 by $f_1^{-1}(v_i) = f^{-1}(v_i)$ for $i = 1'', 3, 4, 5$, $f_1^{-1}(v_{1'}) = f^{-1}(v_{1'}) \setminus C_2$ and $f_1^{-1}(v_2) = f^{-1}(v_2) \cup C_2$. We call this operation *moving w_2 from $U_{1'}$ to U_2* .

$$|f_1^{-1}(v_{1'}) \cup f_1^{-1}(v_{1''})| = |f^{-1}(v_{1'}) \cup f^{-1}(v_{1''})| - |C_2| < |f^{-1}(v_{1'}) \cup f^{-1}(v_{1''})|.$$

That contradicts the choice of f . So we have $N(U_2) \cap U_{1'} = N(U_5) \cap U_{1'} = \{u\}$ for some u . Similarly, $N(U_3) \cap U_{1''} = N(U_4) \cap U_{1''} = \{v\}$ for some v .

Since G is quasi-4-connected, if $\{u, v\}$ is a 2-separator, then $|U_1| = 2$. If $|U_1| \geq 3$, then $\{u, v\}$ is not a 2-separator and there exists $w \in U_1 \setminus \{u, v\}$ such that $w \in N(U_i)$ for some $i = 2, 3, 4, 5$. Without loss of generality, we can assume $w \in N(U_2)$.

Since $N(U_2) \cap U_{1'} = \{u\}$, $w \notin U_{1'}$. If the path P from u to v in T passes through w , then we can move w from $U_{1''}$ to $U_{1'}$, which contradicts $w \notin U_{1'}$. If P does not pass through w , then we can move w from $U_{1''}$ to U_2 , which contradicts the choice of f . ■

From Claim 1, we can let $U_{1'} = \{u_{1'}\}$ and $U_{1''} = \{u_{1''}\}$.

Claim 2. U_2 is at most 1-splittable with respect to $J = \{1', 3, 4, 5\}$ with a possible partition $\{\{1', 5\}, \{3, 4\}\}$.

Proof. U_2 is neither $\{\{1', 3\}, \{4, 5\}\}$ -splittable nor $\{\{1', 4\}, \{3, 5\}\}$ -splittable. Otherwise we can have the $(P_{10})_3$ -minors illustrated in Figs. 9 and 10, respectively. ■




Fig. 10.




Fig. 11.




Fig. 12.

By Proposition 3.4(i), U_2 has a 1-separator $(A_2, B_2; x_2)$ such that $[N(U_{1'}) \cup N(U_5)] \cap U_2 \subseteq A_2$ and $[N(U_3) \cup N(U_4)] \cap U_2 \subseteq B_2$, as we can see in Fig. 11.

Symmetrically, we have the following conclusions (as shown in Fig. 12):

- (i) U_5 is at most 1-splittable with respect to $J = \{1', 2, 3, 4\}$ with the only possible (2×2) -partition $\{\{1', 2\}, \{3, 4\}\}$ and it has a 1-separator $(A_5, B_5; x_5)$ such that $[N(U_{1'}) \cup N(U_2)] \cap U_5 \subseteq A_5$, and $[N(U_3) \cup N(U_4)] \cap U_5 \subseteq B_5$.
- (ii) U_3 is at most 1-splittable with respect to $J = \{1'', 2, 4, 5\}$ with the only possible (2×2) -partition $\{\{1'', 4\}, \{2, 5\}\}$ and it has a 1-separator $(A_3, B_3; x_3)$ such that $[N(U_{1'}) \cup N(U_4)] \cap U_3 \subseteq A_3$, and $[N(U_2) \cup N(U_5)] \cap U_3 \subseteq B_3$.
- (iii) U_4 is at most 1-splittable with respect to $J = \{1'', 2, 3, 5\}$ with the only possible (2×2) -partition $\{\{1'', 3\}, \{2, 5\}\}$ and it has a 1-separator $(A_4, B_4; x_4)$ such that $[N(U_{1'}) \cup N(U_3)] \cap U_4 \subseteq A_4$, and $[N(U_2) \cup N(U_5)] \cap U_4 \subseteq B_4$.

Claim 3. $\{N(u_{1'}) \cap A_2 - \{x_2\}\} \cup \{N(u_{1'}) \cap A_5 - \{x_5\}\} \neq \emptyset$.

Proof. Otherwise, $T = \{u_{1'}, x_2, x_5\}$ is a non-trivial 3-separator of G that separates G with $A_2 \cup A_5 \cup U_{1'}$ as one part. By Lemma 3.3, G is quasi-4-connected; therefore, $A_2 \cup A_5 \cup U_{1'}$ is trivial, but it is not acyclic. ■

Similarly, $\{N(u_{1'}) \cap A_3 - \{x_3\}\} \cup \{N(u_{1'}) \cap A_4 - \{x_4\}\} \neq \emptyset$.




Fig. 13.




Fig. 14. A $(P_{10})_{\bar{3}}$ -minor.

Without loss of generality, we assume that

$$\{N(u_{1'}) \cap A_3 - \{x_3\}\} \neq \emptyset, \quad \{N(u_{1''}) \cap A_2 - \{x_2\}\} \neq \emptyset. \quad (1)$$

Claim 4. U_2 is not $\{\{1'', 5\}, \{3, 4\}\}$ -splittable.

Proof. Otherwise G has a $(P_{10})_{\bar{3}}$ -minor as in Fig. 13 (note that the edge between $U_{1'}$ and U_3 is given by (1)). ■

Symmetrically, U_5 is not $\{\{1'', 2\}, \{3, 4\}\}$ -splittable.

Claim 5. U_2 is at most 0-splittable with respect to $J = \{1', 3, 4, 5\}$.

Proof. By way of contradiction, assume U_2 is not 0-splittable with respect to $J = \{1', 3, 4, 5\}$. By Claim 2, U_2 is $\{\{1', 5\}, \{3, 4\}\}$ -splittable.

Let $\{P_{1',5}, P_{3,4}\}$ be a pair of vertex disjoint paths in U_2 that P_{ij} joins $N(U_i) \cap U_2$ and $N(U_j) \cap U_2$ for $i, j \in \{1', 3, 4, 5\}$.

It is obvious that $P_{3,4}$ must contain the cut vertex x_2 for otherwise A_2 contains a path joining $N(u_{1''})$ and $N(U_5)$. This contradicts Claim 4. Therefore, $N(U_{1'}) \cap (A_2 - x_2) \neq \emptyset$, $N(U_5) \cap (A_2 - x_2) \neq \emptyset$ and both of them are contained in the same component of $A_2 - x_2$, called C_2 , while $N(U_{1''}) \cap (A_2 - x_2)$ is contained in another component of $A_2 - x_2$.

Symmetrically, $A_5 - x_5$ has a component C_5 that contains $N(U_{1'}) \cap (A_5 - x_5)$ and $N(U_2) \cap (A_5 - x_5)$ and is disjoint with $N(U_{1''})$.

Here we have obtained a 3-separator $(H_1, H_2; T)$ with $T = \{u_{1'}, x_2, x_5\}$ as the cut and $H_1 = C_2 \cup C_5 \cup \{u_{1'}, x_2, x_5\}$. Note that neither H_1 nor H_2 is trivial. This contradicts Lemma 3.3. ■

Similarly, U_3 is at most 0-splittable with respect to $J = \{1'', 3, 4, 5\}$.

Final Step:

By Claim 5 and Proposition 3.4(ii), x_2 separates U_2 into four parts $U_2(1')$, $U_2(5)$, $U_2(4)$ and $U_2(3)$ such that $N(U_i) \cap U_2 \subseteq U_2(i)$ for $i \in \{1', 3, 4, 5\}$.

By Claim 3, $N(u_{1''}) \cap A_2 - \{x_2\} \neq \emptyset$. Assume that $N(u_{1''}) \cap A_2 - \{x_2\} \subseteq U_2(1') - x_2$. Then $\{u_{1'}, u_{1''}, x_2\}$ is a 3-separator of G with $U_2(1') \cup U_1$ as a part. Both parts of G separated by $\{u_{1'}, u_{1''}, x_2\}$ contain cycles. This contradicts Lemma 3.3. So, there exists a vertex $v \in U_2(5) \cap N(u_{1''}) - \{x_2\}$ since $A_2 = U_2(1') \cup U_2(5)$.

Similarly, there is a vertex $w \in N(u_{1'}) \cap A_3 - \{x_3\}$, from which we deduce $w \in U_3(4)$. Now we have a $(P_{10})_{\bar{3}}$ -minor as in Fig. 14.

Acknowledgements

The second author was partially supported by the National Security Agency under Grant MSPR-03G-023 and a WV-RCG grant.

References

- [1] K. Appel, W. Haken, Every map is four colorable, Part I: Discharging, *Illinois J. Math.* 21 (1977) 429–490.
- [2] K. Appel, W. Haken, J. Koch, Every map is four colorable, Part II: Reducibility, *Illinois J. Math.* 21 (1977) 491–567.
- [3] K. Appel, W. Haken, Every map is four colorable, *Contemp. Math. AMS* 98 (1989).
- [4] J.A. Bondy, U.S.R. Murty, *Graph Theory with Applications*, Macmillan, London, 1976. Elsevier, New York.
- [5] P.A. Catlin, Double cycle covers and the Petersen graph, *J. Graph Theory* 13 (1989) 465–483.
- [6] H.-J. Lai, *Matroid Theory*, Chinese Higher Education Press, ISBN: 7-04-010563-2, 2002 (in Chinese).
- [7] H.-J. Lai, X. Li, H. Poon, Nowhere-zero 4-flow in regular matroids, *J. Graph Theory* 49 (2005) 196–204.
- [8] R. Diestel, *Graph Theory*, 2nd ed., Springer-Verlag, New York, 1997.
- [9] J. Oxley, *Matroid Theory*, Oxford University Press, New York, 1992.
- [10] N. Robertson, D. Sanders, P.D. Seymour, R. Thomas, The 4-color theorem, *J. Combin. Theory Ser. B* 70 (1) (1997) 2–44.
- [11] N. Robertson, D. Sanders, P.D. Seymour, R. Thomas, Tutte's edge-colouring conjecture, *J. Combin. Theory Ser. B* 70 (1) (1997) 166–183.
- [12] N. Robertson, D. Sanders, P.D. Seymour, R. Thomas, Personal communication.
- [13] R. Thomas, J.M. Thomson, Excluding minors in nonplanar graphs of girth at least five, *Combin. Probab. Comput.* 9 (2000) 573–585.
- [14] W.T. Tutte, On the algebraic theory of graph colorings, *J. Combin. Theory* 1 (1966) 15–20.
- [15] P.N. Walton, D.J.A. Welsh, On the chromatic number of binary matroids, *Mathematika* 27 (1980) 1–9.
- [16] C.-Q. Zhang, *Integer Flows and Cycle Covers of Graphs*, Marcel Dekker, New York, 1997.