

## Brooks' Graph-Coloring Theorem and the Independence Number\*

PAUL A. CATLIN

*Department of Mathematics, Wayne State University, Detroit, Michigan 48202*

*Communicated by C. St. J. A. Nash-Williams*

Received August 4, 1976

Let  $h$  denote the maximum degree of a connected graph  $H$ , and let  $\chi(H)$  denote its chromatic number. Brooks' Theorem asserts that if  $h \geq 3$ , then  $\chi(H) \leq h$ , unless  $H$  is the complete graph  $K_{h+1}$ . We show that when  $H$  is not  $K_{h+1}$ , there is an  $h$ -coloring of  $H$  in which a maximum independent set is monochromatic. We characterize those graphs  $H$  having an  $h$ -coloring in which some color class consists of vertices of degree  $h$  in  $H$ . Again, without any loss of generality, this color class may be assumed to be maximum with respect to the condition that its vertices have degree  $h$ .

### 1. NOTATION

We shall follow the notation of Harary [3]. All graphs in this paper are simple. Let  $V(G)$  denote the vertex set of the graph  $G$ , and let  $E(G)$  denote the edge set. We shall assume that  $E(G)$  is nonempty. Thus, the maximum degree  $\Delta(G)$  of the vertices of  $G$  is at least 1.

For any set  $X$ , we let  $|X|$  denote the cardinality of  $X$ . To simplify notation, we denote the singleton set  $\{x\}$  by  $x$ , so that the union of a set  $S$  and that singleton may be written  $S + x$ .

A *coloring* of  $G$  is a partition of  $V(G)$  into independent subsets, where the partition is unordered and admits null sets. A set  $X \subseteq V(G)$  is *monochromatic* in a coloring of  $G$  if all vertices of  $X$  have the same color. The *chromatic number*  $\chi(G)$  of  $G$  is the minimum possible number of sets in a coloring of  $G$ .

A  $\theta$ -graph is a graph consisting of three distinct arcs, joining the same two vertices and having no other common vertices.

### 2. INTRODUCTION

The basic result in the literature on the problem of coloring a graph  $G$  of specified maximum degree is Brooks' Theorem [2]:

\* This paper is part of the author's Ph.D. thesis, done at the Ohio State University under Professor Neil Robertson.

THEOREM 2.1. *Let  $G$  be a graph with maximum degree  $\Delta(G)$ . We have*

$$\chi(G) \leq \Delta(G) + 1. \quad (2.1)$$

*If  $\Delta(G) = 2$ , then equality holds in (2.1) if and only if  $G$  contains an odd cycle. If  $\Delta(G) \neq 2$ , then equality holds if and only if  $G$  contains a clique  $K_{\Delta(G)+1}$ .*

Note that if  $\Delta(G) = 2$ , an odd cycle of  $G$  is necessarily a connected component of  $G$ . Also, a clique  $K_{\Delta(G)+1}$  is necessarily a component of  $G$ . Such components, which force equality in (2.1), are called  $B_{\Delta(G)}$ -components.

Since each component of a graph can be colored independently, we can assume without loss of generality, that  $G$  is connected.

We give a proof of Brooks' Theorem by induction on  $\Delta(G)$ , and in so doing, we obtain new information. For instance, we show that if  $G$  is not a  $B_{\Delta(G)}$ -component, then there is a coloring of  $G$  in  $\Delta(G)$  colors in which some monochromatic set has  $\beta(G)$  vertices. Also, we characterize those connected graphs  $G$  for which there is a coloring of  $G$  in  $\Delta(G)$  colors such that some monochromatic set consists solely of vertices of degree  $\Delta(G)$ .

Albertson, Bollobás, and Tucker [1] showed first that with two exceptions  $H_1$  and  $H_2$  defined below, every graph  $H$  with  $\Delta(H) = h$  and with no subgraph  $K_h$  has independence number

$$\beta(H) > |V(H)|/h,$$

and they conjectured that such graphs  $H$  have an  $h$ -coloring in which some monochromatic set has more than  $|V(H)|/h$  vertices. Second, they proved this conjecture for graphs that are not regular of degree  $h$ . Theorem 3.2, below, combined with the first result of Albertson, Bollobás, and Tucker shows that this conjecture is true, even for regular graphs.

The two exceptional graphs  $H_1$  and  $H_2$  may be defined as follows: let  $V(H_1)$  be the integers modulo 8, and let  $\{v, w\} \in E(H_1)$  if and only if

$$v - w \equiv 1, 2, 6, \text{ or } 7 \pmod{8}.$$

Let  $V(H_2)$  be the integers modulo 10, and let  $\{v, w\} \in E(H_2)$  if and only if

$$v - w \equiv 1, 4, 5, 6, \text{ or } 9 \pmod{10}.$$

### 3. THE MAIN RESULTS

In this section, we shall consider a connected graph  $H$  with at least one edge. To simplify notation, we denote  $\Delta(H)$  by  $h$ .

A largest independent subset of the set of vertices of degree  $h$  in  $H$  will be called a *superstable set*.

The equivalence of (3.4) and (3.6) of Theorem 3.2 below is Brooks' Theorem (Theorem 2.1).

A *Brooks tree* is any graph  $H$  with  $\Delta(H) = h$  that arises from a tree  $T$  satisfying  $\Delta(T) \leq h$  by the replacement of each vertex of  $T$  with

- (a) an odd cycle if  $h = 3$ ;
- (b) a clique  $K_h$  if  $h \neq 3$ ,

such that if  $x$  and  $y$  are adjacent vertices of  $T$ , then the cycles or cliques substituted for  $x$  and  $y$  are joined by an edge whose removal disconnects  $H$ . Thus,  $K_2$  is the only Brooks tree with  $h = 1$ ; odd arcs with at least 3 edges are the only Brooks trees with  $h = 2$ ; and if  $h \geq 3$ , then a Brooks tree is not a tree.

**THEOREM 3.1.** *Let  $H$  be a connected graph with  $\Delta(H) = h \geq 1$ . The following are equivalent:*

- (3.1)  $H$  is a  $B_h$ -component, or a Brooks tree;
- (3.2) There is no superstable set  $S$  such that  $H - S$  can be colored in  $h - 1$  colors;
- (3.3) There is no independent set  $S$  of vertices of degree  $h$  such that  $H - S$  can be colored in  $h - 1$  colors.

We also have

**THEOREM 3.2.** *Let  $H$  be a connected graph with  $\Delta(H) = h \geq 1$ . The following are equivalent:*

- (3.4)  $H$  is a  $B_h$ -component;
- (3.5) There is no maximum independent set  $S$ , such that  $H - S$  can be colored in  $h - 1$  colors;
- (3.6) There is no  $h$ -coloring of  $H$ .

*Proof of Theorem 3.2 from Theorem 3.1.* For  $\Delta(H) \leq 2$ , the theorem is easily verified. Assume therefore, that  $\Delta(H) \geq 3$ .

We show that if (3.1), (3.2), and (3.3) are equivalent for  $\Delta(H) = h$ , then (3.4), (3.5), and (3.6) are also equivalent for  $\Delta(H) = h$ . Since (3.4) implies (3.6) and (3.6) implies (3.5), it suffices to prove that (3.5) implies (3.4) if (3.1), (3.2), and (3.3) are equivalent.

Adjoin to  $H$  a set  $V$  of  $\sum (h - \deg_H(v))$  vertices disjoint from  $V(H)$ , where the sum runs over all  $v \in V(H)$ . We join each vertex  $v$  of  $H$  to exactly  $h - \deg_H(v)$  vertices of  $V$ , such that no vertex of  $V$  is joined to more than one vertex of  $H$ . Denote the resulting graph  $H'$ . Then,

$$(3.7) \quad H'[V(H)] = H;$$

$$(3.8) \quad \text{Any } v \in V(H) \text{ has degree } h \text{ in } H';$$

$$(3.9) \quad \text{Any } v \in V \text{ has degree } 1 \text{ in } H'.$$

By (3.7) and (3.8), a superstable set  $S$  in  $H'$  is a maximum independent set in  $H$ . Hence, (3.5) for  $H$  implies (3.2) for  $H'$ , whence by (3.1), either  $H'$  is a  $B_h$ -component, or it is a Brooks tree. Since Brooks trees have vertices of degree  $h - 1$ , conditions (3.8), (3.9), and  $h \geq 3$  imply that  $H'$  is not a Brooks tree. Thus,  $H'$  is a  $B_h$ -component, and, therefore, has no vertices of degree 1, whence  $H = H'$ . This proves (3.4), and thus the equivalence of (3.4), (3.5), and (3.6). Hence, Theorem 3.2 follows from Theorem 3.1.

*Proof of Theorem 3.1.* Again, we may suppose that  $h \geq 3$ . Since (3.1) implies (3.3) and (3.3) implies (3.2), it suffices to show that (3.2) implies (3.1).

Suppose inductively that the theorem is true for all graphs  $G$  with  $\Delta(G) < h$ . Then Theorem 3.2 is true for such graphs  $G$ . Let  $H$  be a graph with  $\Delta(H) = h$  such that  $H$  does not satisfy (3.1), and such that for any superstable set  $S$ ,  $H - S$  has no  $(h - 1)$ -coloring. For a given superstable set  $S$ , Theorem 3.2 and

$$\Delta(H - S) \leq h - 1$$

imply that either  $H - S$  can be colored in  $h - 1$  colors, or  $H - S$  has a  $B_{h-1}$ -component. We have already precluded the first possibility. Hence,  $H - S$  has a  $B_{h-1}$ -component. Without loss of generality, we shall choose  $S$  to be a superstable set that minimizes the number of  $B_{h-1}$ -components in  $H - S$ .

Suppose that a vertex  $s \in V(H)$  is in no  $B_{h-1}$ -component in  $H - S$ , regardless of the choice of a superstable set  $S$  that minimizes the number of  $B_{h-1}$ -components in  $H - S$ . Since  $H$  is connected, such a vertex  $s$  exists that is adjacent to a vertex  $v$  lying in a  $B_{h-1}$ -component  $C$  of  $H - S$ , for some such  $S$ . Since the only vertex not in  $C$  that is adjacent to  $v$  lies in  $S$ , we must have  $s \in S$ . Then  $S + v - s$  is a superstable set, and either  $H - (S + v - s)$  has one fewer  $B_{h-1}$ -component than  $H - S$ , contrary to the choice of  $S$ , or  $s$  lies in a  $B_{h-1}$ -component of  $H - (S + v - s)$ , contrary to the choice of  $s$ . Hence, by contradiction, all vertices of  $H$  lie in  $B_{h-1}$ -components of  $H - S$ , for suitable  $S$ .

*Case I.* Suppose that there is no cycle  $P$  in  $H$  with the property that it is not contained in a  $B_{h-1}$ -component of  $H - S$ , regardless of the choice of superstable set  $S$  that minimizes the number of  $B_{h-1}$ -components in  $H - S$ . Observe that if  $h = 3$ , then any even cycle has this property, and any cycle not contained in a clique has this property if  $h > 3$ . Hence, since  $P$  does not exist,

$$\text{If } P' \text{ is a cycle in } H \text{ and if } h = 3, \text{ then } P' \text{ has odd girth} \quad (3.10)$$

and

If  $P'$  is a cycle in  $H$  and  $h \geq 4$ , then  $P'$  is contained in a clique. (3.11)

We see that if  $H$  is a Brooks tree or a  $B_h$ -component, then (3.10) holds if  $h = 3$ , and (3.11) holds if  $h > 3$ . However, we shall prove the converse: i.e., if either (3.10) or (3.11) holds, then  $H$  is either a Brooks tree or a  $B_h$ -component. The main step in proving this converse lies in showing that

(3.12) If  $H$  is not  $K_{h+1}$ , then  $H$  can be partitioned into:

- (i) odd cycles, if  $h = 3$  and (3.10) holds or
- (ii) cliques  $K_h$  if  $h > 3$  and (3.11) holds.

Then it follows from (3.10) or (3.11), from (3.12), and from the definition of Brooks trees that  $H$  must be either  $K_{h+1}$  or a Brooks tree.

Thus, if  $P$  does not exist as described, then (3.10) or (3.11) holds, depending upon the value of  $h$ , and (3.1) follows (which is what we were to prove), provided (3.12) is proven. We shall, therefore, complete Case I by proving (3.12).

We have already shown that each vertex of  $H$  lies in a  $B_{h-1}$ -component of  $H - S$  for some superstable set  $S$  that minimizes the number of  $B_{h-1}$ -components in  $H - S$ . If we show that each vertex of  $H$  lies in exactly one such  $B_{h-1}$ -component, then (3.12) follows.

Suppose, therefore, that some vertex of  $H$  lies in a  $B_{h-1}$ -component  $C_1$  of  $H - S_1$  and in a  $B_{h-1}$ -component  $C_2$  of  $H - S_2$ , for suitable  $S_1$  and  $S_2$ , where  $C_1$  and  $C_2$  are distinct. We shall now derive contradictions with (3.10) and (3.11). If  $h \geq 4$ , then  $C_1$  and  $C_2$  are cliques on  $h$  vertices each. Since they overlap,  $\Delta(H) = h$  forces

$$|V(C_1) \cup V(C_2)| \leq h + 1,$$

and since  $C_1$  and  $C_2$  are distinct, we have equality. Hence,  $H[V(C_1) \cup V(C_2)]$  is either isomorphic to  $K_{h+1}$  or to  $K_{h+1}$  minus an edge. In the first case,  $H$  is  $K_{h+1}$ . In the second case, the cycle  $P'$  of 4 vertices in  $H[V(C_1) \cup V(C_2)]$  containing the two nonadjacent vertices violates (3.11). On the other hand, if  $h = 3$ , then  $C_1$  and  $C_2$  are overlapping odd cycles, and  $\Delta(H) = h < 4$  forces them to overlap in an edge. Then  $C_1 \cup C_2$  contains a  $\theta$ -graph, and hence an even cycle, contrary to (3.10).

*Case II.* Suppose that there is a cycle  $P$  in  $H$  that is not contained in a  $B_{h-1}$ -component of  $H - S$ , where  $S$  is superstable and the number of  $B_{h-1}$ -components of  $H - S$  is minimized. This number of  $B_{h-1}$ -components of  $H - S$  is positive, for otherwise, since the induction hypothesis on  $h$  implies that (3.4) and (3.6) are equivalent for  $H - S$ , we would have  $\chi(H - S) \leq h - 1$ , contrary to (3.2). Since every vertex of  $H$  lies in a  $B_{h-1}$ -component of

$H - S$ , for suitable  $S$ , we can choose  $S = S_0$  so that a  $B_{h-1}$ -component  $C_0$  of  $H - S_0$  contains a vertex of  $P$ . Furthermore, we can assume without loss of generality that  $|V(P) \cap S_0|$  is minimized, with respect to these conditions.

Since the degree of any vertex of  $C_0$  in  $H - S_0$  is  $h - 1$ , and since  $\Delta(H) = h$ , an edge of  $P$  lies in  $E(C_0)$ . Since  $P$  is not contained in  $C_0$ , which is an induced subgraph of  $H$ , an edge of  $P$  lies outside  $E(C_0)$ . Therefore, there is a vertex  $v_1$  of  $V(P) \cap V(C_0)$  having one incident edge of  $P$  in  $E(C_0)$  and the other incident edge of  $P$ , say  $\{v_1, s_1\}$ , outside  $E(C_0)$ . Since  $C_0$  is a component in  $H - S_0$ , we have  $s_1 \in S_0$ .

We define the *path determined by  $P$*  to be the closed path consisting of successive vertices of the cycle  $P$ , where the first and second vertices in the path are  $v_1$  and  $s_1$ , respectively. Denote the vertices of  $P \cap S_0$  by  $s_1, s_2, \dots, s_m$ , so that as one travels along the path determined by  $P$ , one encounters the vertices of  $V(P) \cap S_0$  in the order  $s_1, s_2, \dots, s_m$ . For each  $i \leq m$ , let  $v_i$  be the vertex preceding  $s_i$  in the path determined by  $P$ . Define a sequence  $S_1, S_2, \dots, S_{m-1}$  of sets inductively by

$$S_i = S_{i-1} + v_i - s_i.$$

Since  $S_0$  is superstable, so is  $S_1$ . Since  $S_0$  was chosen to minimize the number of  $B_{h-1}$ -components in  $H - S_0$ , and since  $C_0$  is a  $B_{h-1}$ -component of  $H - S_0$  but not of  $H - S_1$ , it follows that  $s_1$  must be contained in a  $B_{h-1}$ -component  $C_1$  of  $H - S_1$ , and  $S_1$  also minimizes the number of  $B_{h-1}$ -components in  $H - S_1$ . Note that  $C_1$  must contain  $v_2$ , and  $v_2$  is adjacent to exactly one vertex of  $S_1$ , namely,  $s_2$ . By repeating this, one sees inductively that for  $i = 1, 2, \dots, m - 1$ ,  $S_i$  is a superstable set such that  $H - S_i$  has the minimum possible number of  $B_{h-1}$ -components, and in particular,  $H - S_i$  has a  $B_{h-1}$ -component  $C_i$  containing the vertices  $s_i$  and  $v_{i+1}$  of  $P$ . By definition of  $s_m$  and of the path determined by  $P$ ,  $s_m$  is adjacent to a vertex of  $V(C_0)$ . By the minimality of  $|V(P) \cap S_0|$ , which we assumed without loss of generality,  $s_m$  is the first vertex after  $s_1$  along the path to be adjacent to a vertex of  $V(C_0)$ . Otherwise, if  $s_n$  were the first and  $n < m$ , then  $P$  could have been formed by passing from  $s_n$  through  $C_0$  to  $v_1$  directly, instead of proceeding from  $s_n$  to  $C_0$  and  $v_1$  by way of  $s_{n+1}$ . This would violate the minimality of  $|V(P) \cap S_0|$ . Since  $s_m$  is the first vertex after  $s_1$  along the path determined by  $P$  to be adjacent to any vertex of  $C_0$ , the component  $C_0 - v_1$  of  $H - S_1$  is also a component of  $H - S_{m-1}$ . Hence, we can define

$$S_m = S_{m-1} + v_m - s_m,$$

knowing not only that  $S_m$  is superstable and that the number of  $B_{h-1}$ -components of  $H - S_m$  is minimized, but also that  $s_m$  is contained in a

$B_{h-1}$ -component  $C_m$  of  $H - S_m$  whose vertices are precisely  $s_m$  and  $V(C_0) - v_1$ . It must follow from the structure of  $B_{h-1}$ -components that

$$N(s_m) - v_m = N(v_1) - s_1,$$

where  $N(v)$  denotes the set of vertices of  $H$  adjacent to  $v$ .

If  $C_0$  is a cycle of girth at least 5, then  $s_m$  is adjacent to two nonadjacent vertices  $x_1, x_2$  of degree  $h = 3$  that comprise  $N(v) - s$ . Since  $s_m$  is the only vertex in  $S_0$  to which  $x_1$  and  $x_2$  are adjacent,  $S_0 \cup \{x_1, x_2\} - s_m$  is a bigger superstable set than  $S_0$ , contrary to the maximality of  $S_0$ .

If  $C_0$  is a clique  $K_h$ ,  $h \geq 3$ , then  $s_m$  is adjacent to every vertex of  $C_0 - v_1$ . If  $v_1$  and  $s_m$  are adjacent, then  $m = 1$ , and  $V(C_0) + s_m$  induces a clique  $K_{h+1}$  in  $H$ . Since  $H$  is connected,  $K_{h+1}$  is necessarily all of  $H$ , a case excluded since (3.1) is false. Suppose, therefore, that  $s_m$  and  $v_1$  are not adjacent. Let  $x$  be a member of the equal sets  $V(C_m - s_m) = V(C_0 - v_1)$ . Then  $H - (S_0 + x - s_m)$  has fewer  $B_{h-1}$ -components than  $H - S_0$ , and  $S_0 + x - s_m$  is a superstable set. This contradicts the assumption that  $S_0$  minimizes the number of  $B_{h-1}$ -components of  $H - S_0$ .

Thus, whether  $h = 3$  or  $h > 3$ , when  $P$  is assumed to exist, we obtain contradictions, and so the theorem is proved.

#### REFERENCES

1. M. ALBERTSON, B. BOLLOBÁS, AND S. TUCKER, The independence ratio and maximum degree of a graph, in "Proc. Seventh South. Conf. on Comb., Graph Theory, and Computing, 1976," pp. 43-50.
2. R. BROOKS, On colouring the nodes of a network, *Proc. Cambridge Philos. Soc.* **37** (1941), 194-197.
3. F. HARARY, "Graph Theory," Addison-Wesley, Reading, Mass., 1969.