

Krzysztof Ciesielski, Department of Mathematics, West Virginia University,  
Morgantown, WV 26506

Lee Larson<sup>1</sup>, Department of Mathematics, University of Louisville, Louisville,  
KY 40292

## THE PEANO CURVE AND $\mathcal{I}$ -APPROXIMATE DIFFERENTIABILITY

### 1 Preliminaries

A function  $f: \mathbb{R} \rightarrow \mathbb{R}$  is *density continuous* ( $\mathcal{I}$ -density continuous, deep- $\mathcal{I}$ -density continuous) at the point  $x$  if it is continuous at  $x$  when the density topology ( $\mathcal{I}$ -density topology, deep- $\mathcal{I}$ -density topology) is used on both the domain and the range [2, 3, 4, 10]. In [4] it is proved that the first coordinate of the classical Peano area-filling curve is nowhere approximately differentiable, even though it is continuous and density continuous. In this paper we generalize this result by proving in Section 3 that the same function is also  $\mathcal{I}$ -density and deep- $\mathcal{I}$ -density continuous, even though it is nowhere  $\mathcal{I}$ -approximately differentiable. To prove this it is shown that a point  $x$  is a deep- $\mathcal{I}$ -density point of the Baire set  $E$  if, and only if,  $x$  is an  $\mathcal{I}$ -density point of the unique regular open set  $\tilde{E}$  such that the symmetric difference  $E \Delta \tilde{E}$  is of the first category.

In Section 4 we give an example of a bounded  $\mathcal{I}$ -approximately continuous function that is not a derivative.

The notation used throughout this paper is standard. In particular,  $\mathbb{R}$  stands for the set of real numbers and  $\mathbb{N} = \{1, 2, 3, \dots\}$ . For  $A, B \subset \mathbb{R}$  and  $d \in \mathbb{R}$  the complement of  $A$  is denoted by  $A^c$ , the symmetric difference of  $A$  and  $B$  is denoted by  $A \Delta B = (A \cup B) \setminus (A \cap B)$ , while  $B - d = \{x - d \in \mathbb{R}: x \in B\}$  and  $dB = \{dx \in \mathbb{R}: x \in B\}$ . The symbol  $\mathcal{B}$  stands for the family of subsets of  $\mathbb{R}$  which have the Baire property and  $\mathcal{I}$  denotes the ideal of

---

<sup>1</sup>Received support from the University of Louisville Graduate School  
AMS Subject Classification 26A21

first category subsets of  $\mathbb{R}$ . A statement about a subset of  $\mathbb{R}$  is true  $\mathcal{I}$ -a.e. if the set on which it fails to be true is in  $\mathcal{I}$ . An open set  $E \subset \mathbb{R}$  is *regular* if  $E = \text{int}(\text{cl}(E))$ . For a set  $E \in \mathcal{B}$  we denote by  $\tilde{E}$  the only regular open set  $A$  for which  $E \Delta A \in \mathcal{I}$ . The Lebesgue measure of a measurable set  $A$  is denoted by  $m(A)$ .

If  $A \in \mathcal{B}$ , then 0 is an  $\mathcal{I}$ -density point of  $A$  if for every increasing sequence  $\{n_m\}_{m \in \mathbb{N}}$  of natural numbers there exists a subsequence  $\{n_{m_p}\}_{p \in \mathbb{N}}$  such that

$$\lim_{p \rightarrow \infty} \chi_{n_{m_p} A \cap (-1,1)} = \chi_{(-1,1)}, \quad \mathcal{I}\text{-a.e.},$$

or, equivalently, that the set

$$\liminf_{p \rightarrow \infty} n_{m_p} A = \bigcup_{q \in \mathbb{N}} \bigcap_{p \geq q} n_{m_p} A$$

is residual in  $(-1,1)$ . We say that a point  $a$  is an  $\mathcal{I}$ -density point of  $A \in \mathcal{B}$  if 0 is an  $\mathcal{I}$ -density point of  $A - a$ . The set of all  $\mathcal{I}$ -density points of  $A \in \mathcal{B}$  is denoted by  $\Phi_{\mathcal{I}}(A)$ . It is not difficult to see that  $A \Delta \Phi_{\mathcal{I}}(A) \in \mathcal{I}$  for every  $A \in \mathcal{B}$  [10, Theorem 3] and that

$$\Phi_{\mathcal{I}}(A) = \Phi_{\mathcal{I}}(B) \quad \text{for every } A, B \in \mathcal{B} \text{ such that } A \Delta B \in \mathcal{I}. \quad (1)$$

The family of sets

$$\mathcal{T}_{\mathcal{I}} = \{A \in \mathcal{B} : A \subset \Phi_{\mathcal{I}}(A)\}$$

forms a topology on  $\mathbb{R}$  called the  $\mathcal{I}$ -density topology [9, 10].

A point  $a \in \mathbb{R}$  is a *deep- $\mathcal{I}$ -density point* [10] of an  $A \in \mathcal{B}$  if there exists a closed set  $F \subset A \cup \{a\}$  such that  $a$  is an  $\mathcal{I}$ -density point of  $F$ . The set of all deep- $\mathcal{I}$ -density points of  $A \in \mathcal{B}$  is denoted by  $\Phi_{\mathcal{D}}(A)$ . The family of sets

$$\mathcal{T}_{\mathcal{D}} = \{A \in \mathcal{B} : A \subset \Phi_{\mathcal{D}}(A)\}$$

forms a topology on  $\mathbb{R}$  called the *deep- $\mathcal{I}$ -density topology* [5, 10].

A point  $x$  is a *dispersion* ( $\mathcal{I}$ -dispersion, deep- $\mathcal{I}$ -dispersion) point of  $A$  if  $x$  is a density ( $\mathcal{I}$ -density, deep- $\mathcal{I}$ -density) point of  $A^c$ .

A function  $f: \mathbb{R} \rightarrow \mathbb{R}$  is said to be  $\mathcal{I}$ -approximately differentiable at a point  $x$  if there exists a number  $D^{(\mathcal{I})} f(x)$ , called the  $\mathcal{I}$ -approximate derivative of  $f$  at  $x$ , such that for every  $\varepsilon > 0$ ,  $x$  is an  $\mathcal{I}$ -density point of some Baire subset of

$$\left\{ t \in \mathbb{R} : \frac{f(t) - f(x)}{t - x} \in (D^{(\mathcal{I})} f(x) - \varepsilon, D^{(\mathcal{I})} f(x) + \varepsilon) \right\}.$$

(Compare also [6] and [10, Definition 8].)

We also need the following easy fact [1, Lemma 4].

**Lemma 1.** *If  $B = \bigcup_{n \in \mathbb{N}} (a_n, b_n)$  is such that  $\lim_{n \rightarrow \infty} b_n = 0$  and there exists a positive number  $c$  such that*

$$\frac{b_n - a_n}{b_n} > c,$$

*for every  $n \in \mathbb{N}$ , then 0 is not an  $\mathcal{I}$ -dispersion point of  $B$ .*

## 2 Basic Lemmas

We start this section with the following lemma.

**Lemma 2.** *If  $A$  is regular open, then  $\Phi_{\mathcal{I}}(A) = \Phi_{\mathcal{D}}(A)$ .*

*Proof.* The inclusion  $\Phi_{\mathcal{D}}(A) \subset \Phi_{\mathcal{I}}(A)$  is obvious from the definitions.

To prove the converse inclusion let  $x \in \Phi_{\mathcal{I}}(A)$ . For simplicity we assume that  $x = 0$ . We must show that 0 is a deep- $\mathcal{I}$ -density point of  $A$ .

But 0 is an  $\mathcal{I}$ -density point of  $A$  if, and only if, 0 is an  $\mathcal{I}$ -dispersion point of  $A^c$ . Without using the full strength of Theorem 1 from [7] it follows that 0 is an  $\mathcal{I}$ -density point of  $A$  if, and only if,

for every nonempty interval  $(a, b) \subset (-1, 1)$  there exist numbers  $\varepsilon > 0$  and  $n_0 \in \mathbb{N}$  such that for every  $n \geq n_0$  there is a nonempty interval  $(c, d) \subset (a, b)$  with the properties:

$$|d - c| > \varepsilon \quad \text{and} \quad (c, d) \cap n\widetilde{A^c} = \emptyset.$$

Notice that in the above it is enough to consider only intervals  $(a, b)$  with rational endpoints. Moreover, by the regularity of  $A$ ,  $\text{cl}(\widetilde{A^c}) = A^c$ , and so

$$\begin{aligned} (c, d) \cap n\widetilde{A^c} = \emptyset &\iff (c, d) \cap nA^c = \emptyset \\ &\iff \frac{1}{n}(c, d) \cap A^c = \emptyset \\ &\iff \frac{1}{n}(c, d) \subset A. \end{aligned}$$

Hence, we can conclude that 0 is an  $\mathcal{I}$ -density point of  $A$  if, and only if,

( $\star$ ) for every nonempty interval  $I = (a, b) \subset (-1, 1)$  with rational endpoints there exists a number  $\varepsilon_I > 0$  and  $n_I \in \mathbb{N}$  such that for every  $n \geq n_I$  there is a nonempty interval  $(c, d) \subset [c, d] \subset (a, b)$  with the properties:

$$|d - c| > \varepsilon_I \quad \text{and} \quad \frac{1}{n}[c, d] \subset A.$$

Now we can construct a closed set  $E \cup \{0\} \subset A \cup \{0\}$  which satisfies ( $\star$ ). This will finish the proof.

Let  $\{I_k\}_{k \in \mathbb{N}}$  be an enumeration of all nonempty subintervals of  $(-1, 1)$  with rational endpoints and assume that  $n_{I_k} \geq k$  for every  $k \in \mathbb{N}$ . For  $k \in \mathbb{N}$ , let  $E_k$  be a union of intervals  $\frac{1}{n}[c_n, d_n]$  for  $n \geq n_{I_k}$ , where  $[c_n, d_n]$  is a subset of  $I_k$  satisfying ( $\star$ ); i.e., such that

$$|d_n - c_n| > \varepsilon_{I_k} \quad \text{and} \quad \frac{1}{n}[c_n, d_n] \subset A.$$

Let

$$E = \bigcup_{k \in \mathbb{N}} E_k.$$

Notice that

$$E_k \subset \left(-\frac{1}{n_{I_k}}, \frac{1}{n_{I_k}}\right) \subset \left(-\frac{1}{k}, \frac{1}{k}\right)$$

so that  $E_k \setminus \left(-\frac{1}{n}, \frac{1}{n}\right)$  intersects only finitely many closed intervals forming  $E_k$ . In particular,

$$E \setminus \left(-\frac{1}{k}, \frac{1}{k}\right) = \bigcup_{i < k} E_i \setminus \left(-\frac{1}{k}, \frac{1}{k}\right),$$

where the right-hand side intersects only finitely many closed intervals from the collection whose union forms  $E$ . This implies that  $E \cup \{0\}$  is closed. This finishes the proof of Lemma 2.

**Lemma 3.** *Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be such that  $f^{-1}(E) \in \mathcal{I}$  for every  $E \in \mathcal{I}$ . Then  $f$  is deep- $\mathcal{I}$ -density continuous if, and only if,  $f$  is  $\mathcal{I}$ -density continuous.*

*Proof.* It is known that every  $\mathcal{I}$ -density continuous function is deep- $\mathcal{I}$ -density continuous [3, Theorem 4.1(iv)]. To prove the converse implication choose a deep- $\mathcal{I}$ -density continuous function  $f$  satisfying the assumption and

let  $f(x)$  be an  $\mathcal{I}$ -density point of  $E \in \mathcal{B}$  with  $f(x) \in E$ . Then, by (1),  $f(x)$  is an  $\mathcal{I}$ -density point of  $\tilde{E}$  and, by the regularity of  $\tilde{E}$  and Lemma 2,  $f(x)$  is also a deep- $\mathcal{I}$ -density point of  $\tilde{E}$ . Thus,  $x$  is a deep- $\mathcal{I}$ -density point of  $f^{-1}(\tilde{E})$ . Moreover, by the assumption,

$$f^{-1}(\tilde{E}) \Delta f^{-1}(E) = f^{-1}(\tilde{E} \Delta E) \in \mathcal{I}.$$

So, by (1),  $x$  is an  $\mathcal{I}$ -density point of  $f^{-1}(E)$ .

We will need also the following characterization of a deep- $\mathcal{I}$ -density point.

**Lemma 4.** *The following conditions are equivalent:*

- (i)  $x$  is a deep- $\mathcal{I}$ -density point of  $A$ ;
- (ii) given  $s \in (0, 1)$ , there exists  $D_s > 0$  and  $R_s \in (0, 1)$  such that whenever  $0 < D < D_s$  and  $(y - \delta, y + \delta) \subset (x - D, x + D) \setminus \{x\}$  with  $2\delta/D > s$ , then there is an interval  $J \subset (y - \delta, y + \delta) \cap A$  with  $m(J)/2\delta > R_s$ .

Proof. Without any loss of generality it may be assumed that  $x = 0$ . In [11, Theorem (5)] Zajíček proves that 0 is an  $\mathcal{I}$ -density point of  $A$  if, and only if, (ii) is satisfied, where the inclusion  $J \subset (y - \delta, y + \delta) \cap A$  is replaced by

$$[(y - \delta, y + \delta) \cap A] \setminus J \in \mathcal{I}. \quad (2)$$

But we can assume that  $A$  is open, since there is a closed set  $F \subset A \cup \{0\}$  for which 0 is an  $\mathcal{I}$ -density point and then, by Lemma 2, 0 is also an  $\mathcal{I}$ -density point of  $\tilde{F} = \text{int}(F)$ . But then (2) implies that  $J \subset (y - \delta, y + \delta) \cap A$ . Lemma 4 is proved.

### 3 Main Theorem

**Theorem 1.** *There exists a continuous, density continuous and  $\mathcal{I}$ -density continuous function  $f$  which is nowhere approximately and  $\mathcal{I}$ -approximately differentiable.*

Proof. We begin by defining a version of the classical Peano area-filling curve  $P : [0, 1] \rightarrow [0, 1] \times [0, 1]$ . To do this, a sequence of continuous paths  $P_n :$

$[0, 1] \rightarrow [0, 1] \times [0, 1]$  for  $n = 0, 1, \dots$ , are defined which converge uniformly to  $P$ . This definition is facilitated by the following basic construction, which will be referred to as BCP.

Given a square  $[a, b] \times [c, d]$  with one of its diagonals a parametrized constant speed path  $\lambda : [\alpha, \beta] \rightarrow [a, b] \times [c, d]$ , we construct a new path  $\lambda_1 : [\alpha, \beta] \rightarrow [a, b] \times [c, d]$  as shown in Figure 1, where the speed of the new path is constant and three times the speed of  $\lambda$ .




Figure 1: Basic construction BCP.

Using symmetries, BCP can be applied to either of the two diagonals of any square with either path orientation. Also, if  $\|G\|_\infty = \sup_x |G(x)|$ , then it is clear that

$$\|\lambda - \lambda'\|_\infty \leq \sqrt{(b-a)^2 + (d-c)^2} \quad (3)$$

for every  $\lambda' : [\alpha, \beta] \rightarrow [a, b] \times [c, d]$ .

To construct the Peano curve, let  $P_0(t) = (t, t)$  and define  $P_1$  by applying BCP to  $P_0$ . The image of  $P_1$  consists of a diagonal from each of the nine squares

$$\left[ \frac{i}{3}, \frac{i+1}{3} \right] \times \left[ \frac{j}{3}, \frac{j+1}{3} \right], \quad i, j = 0, 1, 2.$$

(See Figure 1 with  $a = c = 0$  and  $b = d = 1$ .) Construct  $P_2$  by applying BCP to each of the diagonals of these squares as shown in Figure 2.

This process can be continued inductively in the obvious way to form the sequence  $P_n$ ,  $n \in \mathbb{N}$ . From (3) it follows that

$$\|P_n - P_m\|_\infty \leq \sqrt{2} 3^{-\min(n,m)}.$$




Figure 2: Construction of  $P_2$ .

This shows that  $P_n$  converges uniformly to  $P$ . It is also easy to see that the image of  $P$  is a dense, compact subset of  $[0, 1] \times [0, 1]$ , so  $P$  is an area-filling curve.

If  $P = (p_1, p_2)$ , where  $p_i : [0, 1] \rightarrow [0, 1]$ ,  $i = 1, 2$  are the coordinate functions for  $P$ , then we claim  $f = p_1$  is a function satisfying the conditions of Theorem 1.

To see this, it might be helpful to see how  $f$  can be defined directly as a uniformly convergent sequence of continuous functions  $f_n : [0, 1] \rightarrow [0, 1]$ , where each  $f_n$  is the first coordinate of  $P_n$ . The first coordinate of BCP can be represented by the construction shown in Figure 3. A similar construction can be done with either diagonal via an obvious reflection. This construction is denoted BCX.

Notice that Figure 3(B) also represents  $f_1 : [0, 1] \rightarrow [0, 1]$ , if we take  $a = \alpha = 0$  and  $b = \beta = 1$ . To form  $f_2$  it is enough to apply BCX to each linear segment of  $f_1$ . Then, apply BCX to each linear segment of  $f_2$  to arrive at  $f_3$ , etc.

Evidently,  $f$  is continuous, as the first coordinate of the continuous function  $P$ . Also, as proved in [4], it is density continuous and nowhere approximately differentiable.

In the rest of the proof, we will need the following easy observations.

The function  $P$  is self-similar in the sense that for every  $n \in \mathbb{N}$  and every  $i = 0, 1, \dots, 9^n - 1$ , there exist  $l(i), r(i) \in \{0, 1, \dots, 3^n - 1\}$  such that the




Figure 3: Basic construction BCX.

image

$$P \left( \left[ \frac{i}{9^n}, \frac{i+1}{9^n} \right] \right) = \left[ \frac{l(i)}{3^n}, \frac{l(i)+1}{3^n} \right] \times \left[ \frac{r(i)}{3^n}, \frac{r(i)+1}{3^n} \right], \quad (4)$$

and the path followed is a scaled and reflected copy of the entire path of  $P$  in  $[0, 1] \times [0, 1]$ . Since  $f$  is the first coordinate of  $P$ , condition (4) implies also that for each integer  $i \in \{0, 1, \dots, 9^n - 1\}$ , there is an integer  $l(i) \in \{0, 1, \dots, 3^n - 1\}$  such that

$$f \left( \left[ \frac{i}{9^n}, \frac{i+1}{9^n} \right] \right) = \left[ \frac{l(i)}{3^n}, \frac{l(i)+1}{3^n} \right]. \quad (5)$$

Also notice the following easy geometrical fact.

For every  $t \in \mathbb{N}$ ,  $t > 1$ , and nonempty interval  $(a, b) \subset [0, 1]$  there are  $i, n \in \mathbb{N}$  such that

$$K = \left[ \frac{i}{t^n}, \frac{i+1}{t^n} \right] \subset (a, b) \quad \text{and} \quad \frac{m(K)}{b-a} \geq \frac{1}{2t} \quad (6)$$



To see this, let  $n$  be the smallest natural number such that

$$1/t^n < (b - a)/2.$$

Thus,  $2/t^{n-1} \geq (b-a)$  and there exists  $i$  such that  $i/t^n \in (a, (b+a)/2)$ . Hence,  $K = [i/t^n, (i+1)/t^n] \subset (a, b)$  and  $m(K)/(b-a) \geq (1/t^n)/(2/t^{n-1}) = 1/2t$ . This finishes the proof of (6).

Notice that (5) implies  $f^{-1}(E)$  is nowhere dense for every nowhere dense set  $E$ . So,

$$f^{-1}(E) \in \mathcal{I} \text{ for every } E \in \mathcal{I}.$$

Thus, by Lemma 3, to show that  $f$  is  $\mathcal{I}$ -density continuous it is enough to prove that  $f$  is deep- $\mathcal{I}$ -density continuous.

Let  $x \in [0, 1]$  and let  $A \subset \mathbb{R} \setminus \{f(x)\}$  be a set such that  $f(x)$  is a deep- $\mathcal{I}$ -density point of  $A$ . It must be shown that  $x$  is a deep- $\mathcal{I}$ -density point of  $f^{-1}(A)$ . This will be done with the aid of Lemma 4.

Let  $s = 1/9^k \in (0, 1)$ . We must find  $D_s > 0$  and  $R_s \in (0, 1)$  such that whenever  $0 < D < D_s$  and an interval  $I \subset (x - D, x + D) \setminus \{x\}$  with  $m(I)/D > s$ , then there is an interval  $J \subset I \cap f^{-1}(A)$  with

$$\frac{m(J)}{m(I)} > R_s. \quad (7)$$

Let  $s' = s/9^3$ . Using Lemma 4 with  $A$  and  $f(x)$ , there exists  $D_{s'} > 0$  and  $R_{s'} = 1/3^l \in (0, 1)$  such that

- whenever  $0 < D' \leq D_{s'}$ , and an interval

$$I' \subset (f(x) - D', f(x) + D') \setminus \{f(x)\}$$

with  $m(I')/D' \geq s'$ , then there is an interval  $J' \subset I' \cap A$  with

$$m(J')/m(I') > R_{s'} \quad (8)$$

Let  $D_s > 0$  be such that

$$|f(x) - f(y)| < D_{s'} \text{ for } |x - y| < D_s \quad (9)$$

and let  $R_s = 1/9^{l+5}$ . Let  $0 < D < D_s$  and choose an interval  $I \subset (x - D, x + D) \setminus \{x\}$  with  $m(I)/D > s$ . We will find an interval  $J \subset I \cap f^{-1}(A)$  with  $m(J)/m(I) > R_s$ .

Assume that  $I \subset (x, x + D)$ . The other case is similar.

Using (6), we can find  $I_0 = [j/9^{n-1}, (j+1)/9^{n-1}] \subset I$  such that

$$\frac{m(I_0)}{m(I)} \geq \frac{1}{18}. \quad (10)$$

Moreover, using (4), it is easy to find  $I_1 = [i/9^n, (i+1)/9^n] \subset I_0$  such that  $f(x) \notin f(I_1)$ . Thus,

$$\frac{m(I_1)}{D} = \frac{1}{9} \frac{m(I_0)}{D} \geq \frac{1}{9} \frac{1}{18} \frac{m(I)}{D} > \frac{s}{9^3} = s'.$$

In particular, there exist  $p = (s')^{-1}$  contiguous intervals  $I^1, I^2, \dots, I^p$  of length  $1/9^n$ , one of which is  $I_1$  and such that  $x \in I^1 \cup I^2 \cup \dots \cup I^p$ .

Define

$$D' = \max\{|f(x) - f(i/9^n)|, |f(x) - f((i+1)/9^n)|\} > 0$$

and  $I' = f(I_1)$ . By (9) we see that  $D' < D_{s'}$  and, by (4),  $f(i/9^n)$  and  $f((i+1)/9^n)$  are the end points of  $I'$  so that  $I' \subset [f(x) - D', f(x) + D'] \setminus \{f(x)\}$ . Moreover, since  $x, i/9^n, (i+1)/9^n \in I^1 \cup I^2 \cup \dots \cup I^p$  then, by (5), we have

$$D' \leq m\left(f\left(\bigcup_{j=1}^p I^j\right)\right) \leq \sum_{j=1}^p m(f(I^j)) = pm(I').$$

Hence,

$$\frac{m(I')}{D'} \geq \frac{m(I')}{pm(I')} = p^{-1} = s'.$$

Thus, by (8), there is an interval  $J' \subset I' \cap A$  such that  $m(J')/m(I') > R_{s'}$ .

Using (6), we can find an interval

$$J'_1 = [j_0/3^m, (j_0+1)/3^m] \subset J'$$

such that  $m(J'_1)/m(J') \geq 1/6 > 1/9$ . Hence,

$$\frac{m(J'_1)}{m(f(I_1))} = \frac{m(J'_1)}{m(I')} = \frac{m(J'_1)}{m(J')} \frac{m(J')}{m(I')} > \frac{1}{9} R_{s'} = \frac{1}{3^{l+2}}$$

and  $J'_1 = [j_0/3^m, (j_0+1)/3^m] \subset f(I_1) = f([i/9^n, (i+1)/9^n])$ . But now condition (4) implies easily that there exists an interval

$$J = [j/9^m, (j+1)/9^m] \subset I_1 = [i/9^n, (i+1)/9^n]$$

such that  $f(J) = J'_1$  and

$$\frac{m(J)}{m(I_1)} > \left(\frac{1}{3^{l+2}}\right)^2 = \frac{1}{9^{l+2}}.$$

Hence, by (10),

$$\frac{m(J)}{m(I)} \geq \frac{m(J)}{18m(I_0)} = \frac{1}{9} \frac{m(J)}{18m(I_1)} > \frac{1}{9^3} \frac{1}{9^{l+2}} = R_s.$$

Condition (7) is proved. This finishes the proof that  $f$  is  $\mathcal{I}$ -density continuous.

To see that  $f$  is not  $\mathcal{I}$ -approximately differentiable at a point  $x \in [0, 1]$  let us do the following construction for each  $n \in \mathbb{N}$ . Choose  $i \in \mathbb{N}$  such that  $x \in [i/9^n, (i+1)/9^n]$ . Then, by (5),  $f([i/9^n, (i+1)/9^n]) = [j/3^n, (j+1)/3^n]$  for some  $j \in \mathbb{N}$ . It is also not difficult to see that condition (4) implies that

$$\begin{aligned} & \left\{ f\left(\left[\frac{9i}{9^{n+1}}, \frac{9i+1}{9^{n+1}}\right]\right), f\left(\left[\frac{9i+8}{9^{n+1}}, \frac{9i+9}{9^{n+1}}\right]\right) \right\} \\ &= \left\{ \left[\frac{3j}{3^{n+1}}, \frac{3j+1}{3^{n+1}}\right], \left[\frac{3j+2}{3^{n+1}}, \frac{3j+3}{3^{n+1}}\right] \right\}. \end{aligned}$$

This implies, in particular, that for every  $y \in [9i/9^{n+1}, (9i+1)/9^{n+1}]$  and  $y' \in [(9i+8)/9^{n+1}, (9i+9)/9^{n+1}]$  we have

$$\frac{|f(y) - f(y')|}{|y - y'|} \geq \frac{1/3^{n+1}}{1/9^n} = 3^{n-1}.$$

Hence, an easy geometrical argument implies that for one of the intervals  $[9i/9^{n+1}, (9i+1)/9^{n+1}]$  or  $[(9i+8)/9^{n+1}, (9i+9)/9^{n+1}]$ , which we denote by  $[a_n, b_n]$ , we have  $x \notin [a_n, b_n]$  and

$$\frac{|f(y) - f(x)|}{|y - x|} \geq 3^{n-1} \quad \text{for every } y \in [a_n, b_n].$$

But, by Lemma 1,  $x$  is not an  $\mathcal{I}$ -dispersion point of  $\bigcup_{n \in \mathbb{N}} [a_n, b_n]$ . Thus, for every  $\mathcal{I}$ -density open set  $U$  containing  $x$ , for every  $\varepsilon > 0$  and  $n \in \mathbb{N}$  there is an  $y \in (x - \varepsilon, x + \varepsilon) \cap U \cap \bigcup_{m > n} [a_m, b_m]$  for which

$$\frac{|f(y) - f(x)|}{|y - x|} \geq 3^n.$$

This implies that  $f$  is not  $\mathcal{I}$ -approximately differentiable. Also notice that the construction of the intervals  $[a_n, b_n]$  given above also implies that  $f$  is not approximately differentiable. This finishes the proof of Theorem 1.

## 4 Derivatives and $\mathcal{I}$ -approximate continuity

In this section we show that the well-known fact that every bounded approximately continuous function is a derivative is not true for the bounded  $\mathcal{I}$ -approximately continuous functions.

**Example 1.** *There exists a bounded  $\mathcal{I}$ -density continuous function which is not a derivative.*

Proof. Let  $P \subset (0, 1]$  be a nowhere dense closed set with positive measure. Choose a sequence  $\{n_k\}_{k \in \mathbb{N}}$  of natural numbers satisfying  $\lim_{k \rightarrow \infty} n_k/n_{k+1} = 0$  and define

$$A = \bigcup_{k \in \mathbb{N}} \frac{1}{n_k} P.$$

Then, by [3, Lemma 2.4], 0 is a deep- $\mathcal{I}$ -dispersion point of  $A$ . Hence, there exists a closed set  $B \cup \{0\} \subset A^c$  such that 0 is an  $\mathcal{I}$ -density point of  $B$ . Moreover, it can be assumed that

$$B = \bigcup_{k \in \mathbb{N}} [a_k, b_k] \cup [c_k, d_k],$$

where  $a_k < b_k < a_{k+1} < 0 < d_{k+1} < c_k < d_k$  [5, 8].

On the other hand, for all  $k \in \mathbb{N}$ ,

$$\frac{m(B^c \cap (0, 1/n_k))}{1/n_k} \geq \frac{m(A \cap (0, 1/n_k))}{1/n_k} > m(P) > 0, \quad (11)$$

so 0 is not a dispersion point of  $B^c$ .

Define the function  $f$  on  $A \cup B$  by

$$f(x) = \begin{cases} 1 & x \in A \\ 0 & x \in B \end{cases}$$

and extend  $f$  on elsewhere in such a way that it is piecewise linear on  $(0, \infty)$  and bounded by 1. Since 0 is an  $\mathcal{I}$ -dispersion point of  $B^c$ , it is apparent that  $f$  is  $\mathcal{I}$ -density continuous. On the other hand,  $f$  cannot be a derivative. To see this, suppose  $F$  is any primitive function for  $f$  and define

$$G(x) = \int_0^x f.$$

Since  $f$  is continuous on  $\mathbb{R} \setminus \{0\}$ , we see that  $F - G$  must be constant on both  $(-\infty, 0)$  and  $(0, \infty)$ . Since both  $F$  and  $G$  are continuous, this implies that  $F - G$  is constant on  $\mathbb{R}$  and therefore  $G$  is differentiable on  $\mathbb{R}$ . But, this is impossible since, by (11),

$$\begin{aligned} D^-G(0) &= 0 < m(P) \\ &\leq \liminf_{k \rightarrow \infty} \frac{m(E \cap (0, 1/n_k))}{1/n_k} \\ &\leq \limsup_{k \rightarrow \infty} \frac{G(1/n_k)}{1/n_k} \leq \overline{D}^+G(0). \end{aligned}$$

## References

- [1] V. Aversa and W. Wilczyński. Homeomorphisms preserving  $\mathcal{I}$ -density points. *Boll. Un. Mat. Ital.*, B(7)1:275–285, 1987.
- [2] Krzysztof Ciesielski and Lee Larson. The space of density continuous functions. *Acta Math. Hung.*, to appear.
- [3] Krzysztof Ciesielski and Lee Larson. Various continuities with the density,  $\mathcal{I}$ -density and ordinary topologies on  $\mathbb{R}$ . *Real Anal. Exchange*, 17(1):183–210, 1991-92.
- [4] Krzysztof Ciesielski, Lee Larson, and Krzysztof Ostaszewski. Differentiability and density continuity. *Real Anal. Exchange*, 15:239–247, 1989–90.
- [5] E. Lazarow. The coarsest topology for  $\mathcal{I}$ -approximately continuous functions. *Comment. Math. Univ. Caroli.*, 27(4):695–704, 1986.
- [6] E. Lazarow and W. Wilczyński.  $\mathcal{I}$ -approximate derivatives. *Rad. Mat.*, 5(1):15–27, 1989.
- [7] Ewa Lazarow. On the Baire class of  $\mathcal{I}$ -approximate derivatives. *Proc. Amer. Math. Soc.*, 100(4):669–674, 1987.
- [8] W. Poreda and E. Wagner-Bojakowska. The topology of  $\mathcal{I}$ -approximately continuous functions. *Rad. Mat.*, 2(2):263–277, 1986.

- [9] W. Poreda, E. Wagner-Bojakowska, and W. Wilczyński. A category analogue of the density topology. *Fund. Math.*, 75:167–173, 1985.
- [10] W. Wilczyński. A category analogue of the density topology, approximate continuity, and the approximate derivative. *Real Anal. Exchange*, 10:241–265, 1984-85.
- [11] L. Zajíček. Alternative definitions of the  $J$ -density topology. *Acta Univ. Carolinae–Mat. et Phys.*, 28(1):57–61, 1987.

*Received July 15, 1991*