

**SAMPLE TEST # 1**

Solve the following exercises. **Show your work.** (No credit will be given for an answer with no supporting work shown.)

**Ex. 1.** Each of the following differential equations is of one of the following form: linear, separable, homogenous, Bernouli, or exact. Solve each of these using appropriate method.

(a)  $y' = \frac{e^{-x} + e^x}{3 + 4y}$ ,  $y(0) = 1$

(b)  $\frac{y}{x} + 6x + (\ln x - 2)\frac{dy}{dx} = 0$ ,  $x > 0$

(c)  $ty' - y = t^2e^{-t}$ ,  $t > 0$

(d)  $\frac{dy}{dx} = \frac{x^2 + 3y^2}{2xy}$

(e)  $x^2y' = y^3 - 2xy$ ,  $x > 0$

(f)  $\frac{dy}{dx} + y = \frac{1}{1 + e^x}$

**Ex. 2.** Draw the direction field for the equation  $\frac{dy}{dx} = 1 + y^2$ . In your drawing show the places where the slopes are 1, 2, and 5.

**Ex. 3.** Without solving, determine the largest interval in which the initial value problem  $(x^2 - x - 6)y' + y \cos x = e^x$ ,  $y(2) = 0$ , has a unique solution.

**Ex. 4.** Apply Euler's method to  $y' = \frac{4 - ty}{1 + y^2}$ ,  $y(0) = -2$ , with step  $h = 0.1$  to estimate  $y(0.1)$ .

**Ex. 5.** A tank with a capacity of 500 gal originally contains 200 gal of wather with 100 lb of salt in solution. Water containing 1 lb of salt per gallon is entering at a rate of 3 gal/min, and the mixture is allowed to flaw out of the tank at the rate of 2 gal/min. Write down an initial value problem (ODE plus initial condition) giving the amount of salt in the tank at any time during the first hour. **Do not solve the equation.** Remember to give the initial condition.