MATH 251.007 Instr. K. Ciesielski Fall 2012

SAMPLE TEST # 4

Solve the following exercises. Show your work.

Ex. 1. Set up the integral formulas, including the limits of the integrations, for the following problems. Do not evaluate the integrals!

- (a) The volume of the solid bounded by $z = x^2 + y^2$, z = 0, x = 0, y = 0, and x + y = 1.
- (b) The mass of the plane lamina bounded by $y = x^2$ and y = 2x + 3, with the density $\delta(x, y) = x^2$.
- (c) The mass of the solid T with the density $\delta(x, y, z) = x^2 + e^z$ bounded by the surfaces: 6x + 2y + z = 12, x = 0, y = 0, and z = 0.

Ex. 2. Evaluate the integrals:

(a)
$$\int_0^1 \int_0^{\pi} \frac{1}{x+1} + \sin y \, dy \, dx =$$

(b)
$$\int_{-2}^{0} \int_{0}^{y} (x+2y^{2}) dx dy =$$

(c)
$$\int \int_R \frac{dy \, dx}{\sqrt{9-x^2-y^2}}$$
, where R is the second quadrant region bounded by $x^2+y^2=4$.

Ex. 3. Find the mass of the solid bounded by the hemisphere $x^2 + y^2 + z^2 \le R^2$, $z \ge 0$, with the density $\delta(x, y, z) = x^2 + y^2 + z^2$.

Ex. 4. Find the mass of the plane lamina bounded by x = 0 and $x = 9 - y^2$ with density $\delta(x, y) = x^2$.

Ex. 5. Evaluate $\int_C xy \, ds$, where C is the parametric curve for which x = 3t, $y = t^4$, and $0 \le t \le 1$.

Ex. 6. Evaluate the integral, where C is the graph of $y = x^3$ from (-1, -1) to (1, 1).

$$\int_C y^2 \, dx + x \, dy =$$

Ex. 7. Determine if the following vector field is conservative. Find potential function for a field, if it is conservative.

(a)
$$\mathbf{F} = (x^3 + \frac{y}{x})\mathbf{i} + (y^2 + \ln x)\mathbf{j}$$

(b)
$$\mathbf{F} = (y\cos x + \ln y)\mathbf{i} + \left(\frac{x}{y} + e^y\right)\mathbf{j}$$

Ex. 8. Find a potential function of the vector field and use the fundamental theorem for line integrals to evaluate

$$\int_{(\pi/2,\pi/2)}^{(\pi,\pi)} (\sin y + y \cos x) \, dx + (\sin x + x \cos y) \, dy =$$

Ex. 9. Apply Green's theorem to evaluate the following integral, where the simple closed curve C is the boundary of the circle $x^2 + y^2 = 1$.

$$\oint_C (\sin x - x^2 y) \, dx + xy^2 \, dy =$$