

MATH 251  
Instr. K. Ciesielski  
Spring 2018

**SAMPLE TEST # 3**

Solve the following exercises. **Show your work.** (No credit will be given for an answer with no supporting work shown.)

**Ex. 1.** Show that the following limit does not exist

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^3 y}{x^4 + 7y^4}$$

Solution:

$$\text{On } x\text{-axis, } y = 0: L_1 = \lim_{x \rightarrow 0} \frac{x^3 \cdot 0}{x^4 + 0} = 0.$$

$$\text{On the line } y = x: L_2 = \lim_{x \rightarrow 0} \frac{x^3 \cdot x}{x^4 + 7x^4} = \lim_{x \rightarrow 0} \frac{x^4}{8x^4} = \frac{1}{8}.$$

Answer: Limit does not exist as  $L_1 \neq L_2$ .

**Ex. 2.** Compute the first order partial derivatives of  $f(x, y, z) = ze^{x^2} \cos y$ .

Solution:

$$\frac{\partial f}{\partial x} = f_x = ze^{x^2} \cos y \cdot 2x = 2xze^{x^2} \cos y$$

$$\frac{\partial f}{\partial y} = f_y = ze^{x^2} (-\sin y) = -ze^{x^2} \sin y$$

$$\frac{\partial f}{\partial z} = f_z = e^{x^2} \cos y$$

**Ex. 3.** Compute all second order partial derivatives of  $g(s, t) = e^{5t} + t \sin(3s)$ .

Solution:

$$g_s = 3t \cos(3s) \quad g_{ss} = -9t \sin(3s) \quad g_{st} = 3 \cos(3s)$$

$$g_t = 5e^{5t} + \sin(3s) \quad g_{ts} = 3 \cos(3s) \quad g_{tt} = 25e^{5t}$$

**Ex. 4.** Find an equation of the plane tangent to the surface  $z = x^2 - 5y^3$  at the point  $P(2, 1, -1)$ .

Solution:

$$z_x = 2x; \quad z_x(P) = 2 \cdot 2 = 4;$$

$$z_y = -15y^2; \quad z_y(P) = -15 \cdot 1^2 = -15;$$

$$\text{Normal vector } \mathbf{n} = \langle z_x(P), z_y(P), -1 \rangle = \langle 4, -15, -1 \rangle.$$

$$\text{Answer: } 4(x - 2) - 15(y - 1) - 1(z + 1) = 0 \quad \text{or} \quad 4x - 15y - z + 6 = 0.$$

**Ex. 5.** Find the absolute maximum and the absolute minimum of the function  $f(x, y) = x^3 - xy$  on the region bounded below by parabola  $y = x^2 - 1$  and above by line  $y = 3$ . You will get credit **only** if **all** critical points are found.

Solution: The curves intersect, when  $x^2 - 1 = 3$ , that is, when  $x = \pm 2$ .

Thus, we need to consider the region above  $x^2 - 1$  and below 3 for  $x$  in the interval  $[-2, 2]$ .

**Region's interior:**  $f_x(x, y) = 3x^2 - y$  and  $f_y(x, y) = -x$ . This leads to system  $3x^2 - y = 0$  and  $-x = 0$ , with only solution  $(x, y) = (0, 0)$ . This point belongs to the region. This is our first critical point.

**Lower boundary:**  $y = x^2 - 1$  and  $-2 \leq x \leq 2$ . Then

$$g(x) = f(x, x^2 - 1) = x^3 - x(x^2 - 1) = x \text{ and } g'(x) = 1 \text{ is never } 0.$$

So, there are no true critical points, but we need to consider the endpoints of  $g$ ,  $x = \pm 2$ .

This give us the critical points  $(x, y) = (\pm 2, 3)$ .

**Upper boundary:**  $y = 3$  and  $-2 \leq x \leq 2$ . Then

$$g(x) = f(x, 3) = x^3 - 3x \text{ and } g'(x) = 3x^2 - 3, \text{ which is } 0 \text{ when } x = \pm 1 \in [-2, 2].$$

This give us the critical points  $(x, y) = (\pm 1, 3)$ . (Plus the end points  $(x, y) = (\pm 2, 3)$ , considered above.)

**Checking the critical points:**  $f(0, 0) = 0$ ;

$$f(2, 3) = 2^3 - 6 = 2; f(-2, 3) = (-2)^3 + 6 = -2;$$

$$f(1, 3) = 1^3 - 3 = -2; f(-1, 3) = (-1)^3 + 3 = 2;$$

**Answer:**  $f$  has the absolute maximum value 2, at points  $(2, 3)$  and  $(-1, 3)$ .

$f$  has the absolute minimum value  $-2$ , at points  $(-2, 3)$  and  $(1, 3)$ .

**Ex. 6.** Find the volume of the solid bounded above by the surface  $z = 28xy$ , bounded below by  $xy$ -plane, and which is above the region bounded by  $y = x^6$  and  $y = x$ .

Solution: The curves intersect, when  $x^6 = x$ , that is, when  $x = 0$  and  $x = 1$ .

Thus, we need to find an integral above  $x^6$  and below  $x$ , on the interval  $[0, 1]$ :

$$\int_0^1 \int_{x^6}^x 28xy \, dy \, dx = \int_0^1 [14xy^2]_{y=x^6}^{y=x} \, dx = \int_0^1 [14x^3 - 14x^{13}]_{y=x^6}^{y=x} \, dx = \left[ \frac{14}{4}x^4 - x^{14} \right]_{x=0}^{x=1} = 2.5$$

**Ex. 7.** Evaluate  $\int_0^1 \int_0^x 4e^{x^2} dy dx$

Solution:  $\int_0^1 \int_0^x 4e^{x^2} dy dx = \int_0^1 [4e^{x^2} y]_{y=0}^{y=x} dx = \int_0^1 4(e^{x^2} x - e^{x^2} 0) dx = \int_0^1 4e^{x^2} x dx$

Using substitution  $v = x^2$ , we obtain that it is equal

$$\left[ 2e^{x^2} \right]_{x=0}^{x=1} = 2(e^1 - e^0) = 2(e - 1).$$

**Ex. 8.** Find the point on the cone  $z = \sqrt{x^2 + y^2}$  which is the closest to the point  $(4, -8, 0)$ .

Solution: Distance of  $(x, y, z)$  on the surface from  $(4, -8, 0)$  is  $\sqrt{(x-4)^2 + (y+8)^2 + (z-0)^2}$ .

Since  $z^2 = x^2 + y^2$ , this is equal to

$$f(x, y) = \sqrt{(x-4)^2 + (y+8)^2 + (x^2 + y^2)}.$$

$$f_x(x, y) = \frac{2(x-4)+2x}{2\sqrt{(x-4)^2+(y+8)^2+(x^2+y^2)}} \text{ and } f_y(x, y) = \frac{2(y+8)+2y}{2\sqrt{(x-4)^2+(y+8)^2+(x^2+y^2)}}.$$

$$f_x = 0 \text{ when } 2(x-4) + 2x = 0, \text{ that is, } 4x - 8 = 0, \text{ so } x = 2.$$

$$f_y = 0 \text{ when } 2(y+8) + 2y, \text{ that is, } 4y + 16 = 0, \text{ so } y = -4.$$

This gives critical point  $(2, -4)$ . Since these are the coordinates of a point on the cone, we get  $z = \sqrt{2^2 + (-4)^2} = \sqrt{20}$ .

Answer: Point  $(2, -4, \sqrt{20})$ .

**Ex. 9.** Find the directional derivative of  $f(x, y) = 10e^y \sin x$  at the point  $P(\pi/4, 0)$  in the direction of the vector  $\mathbf{v} = 4\mathbf{i} - 3\mathbf{j}$ .

Solution: The unit vector in the direction of  $\mathbf{v}$  is equal

$$\mathbf{u} = \frac{1}{|\mathbf{v}|} \mathbf{v} = \frac{1}{\sqrt{4^2 + (-3)^2}} \mathbf{v} = \frac{1}{5} \langle 4, -3 \rangle = \langle .8, -.6 \rangle.$$

$$f_x(x, y) = 10e^y \cos x; f_x(P) = 10e^0 \cos(\pi/4) = 5\sqrt{2}.$$

$$f_y(x, y) = 10e^y \sin x; f_y(P) = 10e^0 \sin(\pi/4) = 5\sqrt{2}.$$

$$\nabla f(P) = \langle f_x(P), f_y(P) \rangle = \langle \sqrt{2}/2, \sqrt{2}/2 \rangle.$$

$$D_{\mathbf{u}} f(P) = \nabla f(P) \cdot \mathbf{u} = \langle 5\sqrt{2}, 5\sqrt{2} \rangle \cdot \langle .8, -.6 \rangle = (5\sqrt{2})(.8) + (5\sqrt{2})(-.6) = \sqrt{2}.$$

**Ex. 10.** Find the gradient of  $g(x, y, z) = x^2 + e^{yz} + \cos(x + 2y)$ .

Solution:  $\nabla g(x, y, z) = \langle g_x, g_y, g_z \rangle = \langle 2x - \sin(x + 2y), ze^{yz} - 2 \sin(x + 2y), ye^{yz} \rangle.$