

MATH 251  
Instr. K. Ciesielski  
Spring 2020

**SAMPLE FINAL TEST**  
(longer than the actual Final Test)

Solve the following exercises. **Show your work.**

**Ex. 1. ST #1 Ex 3:** Find the determinant of the matrix. Each time you expand the the matrix, you **must** expand it over a row or column that has the largest number of zeros. If necessary, use the row (or column) reduction method to create additional zeros.

$$A = \begin{bmatrix} -1 & 2 & 0 & 0 \\ 1 & -1 & 1 & -1 \\ 1 & 2 & 0 & 1 \\ 0 & 3 & 1 & 2 \end{bmatrix}$$

**Ex. 2. ST #1 Ex 4:** Find the inverse matrix of

$$A = \begin{bmatrix} 1 & 0 & 1 \\ -1 & 1 & 2 \\ 0 & 1 & -1 \end{bmatrix}$$

**Ex. 3. ST #1 Ex 6:** Let  $\mathbf{a} = \langle 0, 1, 2 \rangle$ ,  $\mathbf{b} = \langle -1, 0, 7 \rangle$ , and  $\mathbf{c} = \langle 2, 3, -1 \rangle$ . Evaluate:  $2\mathbf{a} - \mathbf{b} + \mathbf{c}$ ,  $|\mathbf{c}|$ , and  $(\mathbf{a} \cdot \mathbf{b})(\mathbf{b} \times \mathbf{c})$ . (Do not confuse vectors with numbers. No partial credit for solutions with such errors.)

**Ex. 4. ST #2 Ex 1:** Find a vector equation of the line that passes through the point  $P(11, 13, -7)$  and is perpendicular to the plane with the equation:  $x - 2z = 17$ .

**Ex. 5. ST #2 Ex 7:** Let  $\mathbf{v}(t) = \mathbf{i}(t + e)^{-1} + \mathbf{k} t^3$  be a velocity of a particle. Find the acceleration vector  $\mathbf{a}(t)$  of the particle and its position vector  $\mathbf{r}(t)$ , where its initial position was  $\mathbf{r}(0) = 3\mathbf{i}$ .

**Ex. 6. ST #2 Ex 10:** Sketch and fully describe the domain of the following function, including the name of the surface representing the domain's boundary:  $f(x, y, z) = \ln(25 - 4x^2 - 9y^2 - z^2)$ .

**Ex. 7. ST #3 Ex. 2:** Compute the first order partial derivatives of  $f(x, y, z) = ze^{x^2} \cos y$ .

**Ex. 8. ST #3 Ex. 3:** Compute all second order partial derivatives of  $g(s, t) = e^{5t} + t \sin(3s)$ .

**Ex. 9. ST #3 Ex. 4:** Find an equation of the plane tangent to the surface  $z = x^2 - 5y^3$  at the point  $P(2, 1, -1)$ .

**Ex. 10. ST #3 Ex. 8:** Find the point on the cone  $z = \sqrt{x^2 + y^2}$  which is the closest to the point  $(4, -8, 0)$ .

**Ex. 11. ST #3 Ex. 5:** Find the absolute maximum and the absolute minimum of the function  $f(x, y) = x^3 - xy$  on the region bounded below by parabola  $y = x^2 - 1$  and above by line  $y = 3$ . You will get credit **only** if **all** critical points are found.

**Ex. 12. ST #4 Ex. 1(a)&(c):** Set up the integral formulas, **including the limits of the integrations**, for the following problems. *Do not evaluate the integrals!*

- (a) The volume of the solid bounded by  $z = x^2 + y^2$ ,  $z = 0$ ,  $x = 0$ ,  $y = 0$ , and  $x + y = 1$ .
- (c) The mass of the solid  $T$  with the density  $\delta(x, y, z) = x^2 + e^z$  bounded by the surfaces:  $6x + 2y + z = 12$ ,  $x = 0$ ,  $y = 0$ , and  $z = 0$ .

**Ex. 13. ST #4 Ex. 2:** Evaluate the integrals:

(a)  $\int_0^1 \int_0^\pi \frac{1}{x+1} + \sin y \, dy \, dx =$

(b)  $\int_{-2}^0 \int_0^y (x + 2y^2) \, dx \, dy =$

(c)  $\int \int_R \frac{dy \, dx}{\sqrt{9 - x^2 - y^2}}$ , where  $R$  is the *second quadrant* region bounded by  $x^2 + y^2 = 4$ .

**Ex. 14. ST #4 Ex. 3:** Find the mass of the solid bounded by the hemisphere  $x^2 + y^2 + z^2 \leq R^2$ ,  $z \geq 0$ , with the density  $\delta(x, y, z) = x^2 + y^2 + z^2$ .

**Ex. 15. ST #4 Ex. 4:** Find the mass of the plane lamina bounded by  $x = 0$  and  $x = 9 - y^2$  with density  $\delta(x, y) = y^2$ .

**Ex. 16. ST #4 Ex. 6:** Evaluate the integral, where  $C$  is the graph of  $y = x^3$  from  $(-1, -1)$  to  $(1, 1)$

$$\int_C y^2 \, dx + x \, dy =$$

**Ex. 17. ST #4 Ex. 8:** Find a potential function of the vector field and use the fundamental theorem for line integrals to evaluate

$$\int_{(\pi/2, \pi/2)}^{(\pi, \pi)} (\sin y + y \cos x) \, dx + (\sin x + x \cos y) \, dy =$$

**Ex. 18. ST #4 Ex. 9: Apply Green's theorem** to evaluate the following integral, where the simple closed curve  $C$ , with counter clockwise direction, is the boundary of the circle  $x^2 + y^2 = 1$ .

$$\oint_C (\sin x - x^2 y) \, dx + xy^2 \, dy =$$