Directions: You may work to solve these problems in groups, but all written work must be your own. Unless the problem indicates otherwise, all problems require some justification; a correct answer without supporting reasoning is not sufficient. Submissions must be stapled. See "Guidelines and advice" on the course webpage for more information.

- 1. Proofs involving sets. Prove the following.
 - (a) $\{6n: n \in \mathbb{Z}\} = \{2n: n \in \mathbb{Z}\} \cap \{3n: n \in \mathbb{Z}\}.$
 - (b) $\{9^n: n \in \mathbb{Q}\} = \{3^n: n \in \mathbb{Q}\}$
 - (c) If A and B be sets, then $A \subseteq B$ if and only if $A \cap B = A$.
 - (d) $\bigcap_{x \in \mathbb{R}} [3 x^2, 5 + x^2] = [3, 5].$
- 2. Suppose $B \neq \emptyset$ and $A \times B \subseteq B \times C$. Prove that $A \subseteq C$.
- 3. Each of the following is true or false. Decide which is the case and prove or disprove accordingly, using any method.
 - (a) For every natural number n, the integer $n^2 + 17n + 17$ is prime.
 - (b) If $a, b, c \in \mathbb{N}$ and ab, bc, and ac all have the same parity, then a, b, and c all have the same parity.
 - (c) If A and B are finite sets, then $|A \cup B| = |A| + |B|$.
 - (d) If A and B are sets, then $\mathcal{P}(A) \cap \mathcal{P}(B) = \mathcal{P}(A \cap B)$.
 - (e) If A and B are sets, then $\mathcal{P}(A) \cup \mathcal{P}(B) = \mathcal{P}(A \cup B)$.
 - (f) If p and q are prime numbers for which p < q, then $2p + q^2$ is odd.
- 4. Suppose that $\alpha \in \mathbb{R}$ and $0 < \alpha < 1$. A magical cake has icing on one side. A baker cuts the cake to make a slice with center angle $\alpha \cdot 2\pi$ (radians), flips the slice over (so the piece has icing face-down), and the slice magically reattaches to the rest of the cake. The baker continues to make wedge slices with center angle $\alpha \cdot 2\pi$, proceeding counter-clockwise around the cake with each subsequent slice starting where the previous slice ended. For which $\alpha \in \mathbb{R}$ will this process eventually lead to the cake again having all its icing on the same side (up or down)? For which $\alpha \in \mathbb{R}$ will this process lead to the cake having all its icing facing down?